

Toward an understanding of entanglement for generalized n -qubit W-states

Levon Tamaryan

Physics Department, Yerevan State University, Yerevan, 375025, Armenia

Hungsoo Kim

The Institute of Basic Science, Kyungnam University, Masan, 631-701, Korea

Eylee Jung, Mi-Ra Hwang, DaeKil Park

Department of Physics, Kyungnam University, Masan, Korea

Sayatnova Tamaryan

Theory Department, Yerevan Physics Institute, Yerevan, Armenia

Abstract

We solve stationarity equations of the geometric measure of entanglement for multi-qubit W-type states. In this way we compute analytically the maximal overlap of one-parameter n -qubit and two-parameter four-qubit W-type states and their nearest product states. Possible extensions to arbitrary W-type states and geometrical interpretations of these results are discussed in detail.

I. INTRODUCTION

Entanglement of quantum states [1] plays an important role in quantum information, computation and communication (QICC). It is a genuine physical resource for the teleportation process [2, 3] and makes it possible that the quantum computer outperforms classical one [4, 5]. It also plays a crucial role in quantum cryptographic schemes [6, 7]. These phenomena have provided the basis for the development of modern quantum information science.

Quantum entanglement is a rich field of research. A better understanding of quantum entanglement, of ways it is characterized, created, detected, stored and manipulated, is theoretically the most basic task of the current QICC research. In bipartite case entanglement is relatively well understood, while in multipartite case even quantifying entanglement of pure states is a great challenge.

The geometric measure of entanglement can be considered as one of the most reliable quantifiers of multipartite entanglement [8, 9, 10]. It depends on P_{max} , the maximal overlap of a given state with the nearest product state, and is defined by the formula $E_g(\psi) = 1 - P_{max}$ [10]. The same overlap P_{max} , known also as the injective tensor norm of ψ [11], is the maximal probability of success in the Grover's search algorithm [12] when the state ψ is used as an input state. This relationship between the success probability of the quantum search algorithm and the amount of entanglement of the input state allows oneself to define an operational entanglement measure known as Groverian entanglement [13, 14].

The maximal overlap P_{max} is a useful quantity and has several practical applications. It has been used to study quantum phase transitions in spin models [15, 16] and to quantify the distinguishability of multipartite states by local means [17]. Moreover, P_{max} exhibits interesting connections with entanglement witnesses and can be efficiently estimated in experiments [18]. Recently, it has been shown that the maximal overlap is the largest coefficient of the generalized Schmidt decomposition and the nearest product state uniquely defines the factorizable basis of the decomposition [19, 20].

In spite of its usefulness one obstacle to use P_{max} fully in quantum information theories is the fact that it is difficult to compute it analytically for generic states. The usual maximization method generates a system of nonlinear equations [10]. Thus, it is important to develop a technique for the computation of P_{max} [21, 22, 23, 24, 25].

Theorem I of Ref.[21] enables us to compute P_{max} for n -qubit pure states by making use of $(n-1)$ -qubit reduced states. In the case of three-qubit states the theorem effectively changes the nonlinear eigenvalue equations into the linear form. Owing to this essential simplification P_{max} for the generalized three-qubit W-state [26, 27] was computed analytically in Ref.[28]. Furthermore, in Ref.[29] P_{max} was found for three-qubit quadrilateral states with an elegant geometric interpretation. More recently, based on the analytical results of Ref.[28, 29] and the classification of Ref.[30], P_{max} for various types of three-qubit states was computed analytically and expressed in terms of local unitary(LU) invariants [31].

In general, the calculation of the multi-partite entanglement is confronted with great difficulties. Furthermore, even if we know explicit expressions of entanglement measure, the separation of the applicable domains is also a nontrivial task [29]. Therefore, there is a good reason to consider first some solvable cases that allow analytic solutions and clear separations of the validity domains. Later, these results could be extended, either analytically or numerically, for a wider class of multi-qubit states. In the light of these ideas we consider one- and two-parametric n -qubit W-type states with $n \geq 4$ in this paper.

The paper is organized as follows. In Sec. II we clarify our tasks and notations. In Sec. III we review the calculational tool introduced in Ref.[21, 28, 29] and explain how the Lagrange multiplier method gives simple solution to the one-parameter cases. This method is used Sec. IV for the derivation of P_{max} for one-parameter W-states in 4-qubit, 5-qubit and 6-qubit systems. In this section the analytical results are compared with numerical data. In Sec. V based on the analytical results of the previous section we compute P_{max} for an one-parameter W-state in arbitrary n -qubit system. In Sec. VI we derive P_{max} for two-parameter W-states in 4-qubit system by adopting the usual maximization technique. In Sec.VII we analyze two-parameter results by considering several particular cases. In Sec. VIII we discuss the possibility of extensions of the results to arbitrary W states and the existence of a geometrical interpretation.

II. SUMMARY OF TASKS

Let $|\psi\rangle$ be a pure state of an n -party system $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_n$, where the dimensions of the individual state spaces $\mathcal{H}_k$ are finite but otherwise arbitrary. The maximal overlap of

$|\psi\rangle$ is given by

$$P_{max}(\psi) \equiv \max_{|q_1\rangle\cdots|q_n\rangle} |\langle q_1|\langle q_2|\cdots\langle q_n|\psi\rangle|^2, \quad (1)$$

where the maximum is taken over all single-system normalized state vectors $|q_k\rangle \in \mathcal{H}_k$, and it is understood that $|\psi\rangle$ is normalized.

Let us consider now n -qubit W-type state

$$|W_n\rangle = a_1|10\cdots 0\rangle + a_2|010\cdots 0\rangle + \cdots + a_n|0\cdots 01\rangle, \quad (2)$$

where the labels within each ket refer to qubits $1, 2, \cdots, n$ in that order.

In this paper we will compute analytically P_{max} in the following two cases:

- 1) for the one-parametric $|W_n\rangle$ when $a_1 = \cdots = a_{n-1} \equiv a$ and $a_n \equiv q$
- 2) for the two-parametric $|W_4\rangle$ when $a_1 = a$, $a_2 = b$, $a_3 = a_4 = q$.

To ensure the calculational validity we use the result of [14], which has shown that $P_{max} = (1 - 1/n)^{n-1}$ when $a_1 = a_2 = \cdots = a_n$. Thus, the final results of the one-parametric case should agree with the following:

- If $a = q = 1/\sqrt{n}$, then P_{max} should be equal to $(1 - 1/n)^{n-1}$.
- If $q = 0$, then $|W_n\rangle$ becomes $|W_{n-1}\rangle \otimes |0\rangle$ and, as a result, P_{max} should be equal to $(1 - 1/(n-1))^{n-2}$.

For the two-parametric case $P_{max}(W_4)$ should have a correct limit when either a or b vanishes. At $a = 0$ we have $|W_4\rangle = |0\rangle \otimes |W_3\rangle$ and thus the maximal overlap should be expressed in terms of the circumradius of the isosceles triangle with sides b, q, q [28].

III. CALCULATION TOOL

For a pure state of two qubits P_{max} is given by

$$P_{max} = \frac{1}{2} \left[1 + \sqrt{1 - 4 \det \rho^A} \right], \quad (3)$$

where ρ^A is reduced density matrix, i.e. $\text{Tr}_B \rho^{AB}$. Therefore, the Bell (and their LU-equivalent) states have the minimal ($P_{max} = 1/2$) while product states have the maximal ($P_{max} = 1$) overlap.

The explicit dependence of P_{max} on state parameters for the generalized three-qubit W-state

$$|W_3\rangle = a_1|100\rangle + a_2|010\rangle + a_3|001\rangle \quad (4)$$

was computed in Ref.[28]. In order to express explicitly $P_{max}(W_3)$ in terms of state parameters, we define a set $\{\alpha, \beta, \gamma\}$ as the set $\{a_1, a_2, a_3\}$ in decreasing order. Then P_{max} for the generalized W-state can be expressed in a form

$$P_{max}(W_3) = \begin{cases} 4R_W^2 & \text{when } \alpha^2 \leq \beta^2 + \gamma^2 \\ \alpha^2 & \text{when } \alpha^2 \geq \beta^2 + \gamma^2 \end{cases} \quad (5)$$

where R_W is the circumradius of the triangle with sides a_1, a_2, a_3 . Similar calculation procedure can be applied to the 3-qubit quadrilateral state. It has been shown in Ref.[29] that for this case P_{max} is expressed in terms of the circumradius of a convex quadrangle. These two separate results strongly suggest that P_{max} for an arbitrary pure state has its own geometrical meaning. If we are able to know this meaning completely, then our understanding on the multipartite entanglement would be greatly enhanced.

Now, we briefly review how to derive the analytic result (5) because it plays crucial role in next two sections. In Ref.[28] P_{max} for 3-qubit state is expressed as

$$P_{max} = \frac{1}{4} \max_{|\vec{s}_1|=|\vec{s}_2|=1} [1 + \vec{s}_1 \cdot \vec{r}_1 + \vec{s}_2 \cdot \vec{r}_2 + g_{ij}s_{1i}s_{2j}] \quad (6)$$

where $\vec{s}_1$ and $\vec{s}_2$ are Bloch vectors of the single-qubit states. In Eq.(6) $\vec{r}_1 = Tr[\rho^A \vec{\sigma}]$, $\vec{r}_2 = Tr[\rho^B \vec{\sigma}]$ and $g_{ij} = Tr[\rho^{AB} \sigma_i \otimes \sigma_j]$, where ρ^A , ρ^B and ρ^{AB} are appropriate partial traces of $\rho^{ABC} \equiv |W_3\rangle\langle W_3|$ and σ_i are usual Pauli matrices. The explicit expressions of $\vec{r}_1$, $\vec{r}_2$ and g_{ij} are given in Ref.[28]. Due to maximization over $\vec{s}_1$ and $\vec{s}_2$ in Eq.(6) we can compute $\vec{s}_1$ and $\vec{s}_2$ by solving the Lagrange multiplier equations

$$\vec{r}_1 + g\vec{s}_2 = \lambda_1 \vec{s}_1, \quad \vec{r}_2 + g^T \vec{s}_1 = \lambda_2 \vec{s}_2, \quad (7)$$

where λ_1 and λ_2 are Lagrange multiplier constants. Now, we let $s_{1y} = s_{2y} = 0$ for simplicity, because those give only irrelevant overall phase factor to $\langle q_1 | \langle q_2 | \langle q_3 | W_3 \rangle$. After eliminating the Lagrange multiplier constants, one can show that Eq.(7) reduces to two equations. Examining these two remaining equations, one can show that $\vec{s}_1$ and $\vec{s}_2$ have a following relation to each other:

$$\vec{s}_1(a_1, a_2, a_3) = \vec{s}_2(a_2, a_1, a_3). \quad (8)$$

Using this relation, one can combine these two equations into single one expressed in terms of solely s_{1z} in a final form

$$\frac{\sqrt{1 - s_{1z}^2(a_1, a_2, a_3)}}{s_{1z}(a_1, a_2, a_3)} = \frac{\omega \sqrt{1 - s_{1z}^2(a_2, a_1, a_3)}}{r_1 - r_3 s_{1z}(a_2, a_1, a_3)} \quad (9)$$

where $r_1 = a_2^2 + a_3^2 - a_1^2$, $r_2 = a_1^2 + a_3^2 - a_2^2$, $r_3 = a_1^2 + a_2^2 - a_3^2$ and $\omega = 2a_1a_2$. Defining $a_1 = a_2 \equiv a$ and $a_3 \equiv q$ again, one can solve Eq.(9) easily in a form

$$\begin{aligned} s_{1z} = s_{2z} &= \frac{r_1}{\omega + r_3} = \frac{q^2}{4a^2 - q^2} \\ s_{1x} = s_{2x} &= \sqrt{1 - s_{1z}^2} = \frac{2\sqrt{2}a}{4a^2 - q^2} \sqrt{2a^2 - q^2}. \end{aligned} \quad (10)$$

Inserting Eq.(10) into Eq.(6), one can compute P_{max} for $|W_3\rangle$ with $a_1 = a_2 = a$ and $s_3 = q$, whose final expression is simply

$$P_{max} = \frac{(1 - q^2)^2}{2 - 3q^2}. \quad (11)$$

Eq.(11) is consistent with Eq.(5) when $q^2 \leq 2a^2$. When $q = 0$, Eq.(11) gives $P_{max} = 1/2$ which corresponds to that of 2-qubit EPR state. When $q = 1/\sqrt{3}$, Eq.(11) gives $P_{max} = 4/9$, which is also consistent with the result of Ref.[14].

IV. FOUR, FIVE AND SIX QUBIT W-TYPE STATES: ONE-PARAMETRIC CASES

The method described in the previous section may enable us to compute P_{max} of four-qubit W-type states. For the case of arbitrary four-qubit systems P_{max} can be represented in a form

$$\begin{aligned} P_{max} &= \frac{1}{8} \max_{|\vec{s}_1|=|\vec{s}_2|=|\vec{s}_3|=1} \left[1 + \vec{s}_1 \cdot \vec{r}_1 + \vec{s}_2 \cdot \vec{r}_2 + \vec{s}_3 \cdot \vec{r}_3 \right. \\ &\quad \left. + s_{1i}s_{2j}g_{ij}^{(3)} + s_{1i}s_{3j}g_{ij}^{(2)} + s_{2i}s_{3j}g_{ij}^{(1)} + s_{i1}s_{2j}s_{3k}h_{ijk} \right], \end{aligned} \quad (12)$$

where

$$\begin{aligned} \vec{r}_1 &= Tr[\rho^A \vec{\sigma}], \quad \vec{r}_2 = Tr[\rho^B \vec{\sigma}], \quad \vec{r}_3 = Tr[\rho^C \vec{\sigma}], \\ g_{ij}^{(3)} &= Tr[\rho^{AB} \sigma_i \otimes \sigma_j], \quad g_{ij}^{(2)} = Tr[\rho^{AC} \sigma_i \otimes \sigma_j], \quad g_{ij}^{(1)} = Tr[\rho^{BC} \sigma_i \otimes \sigma_j] \\ h_{ijk} &= Tr[\rho^{ABC} \sigma_i \otimes \sigma_j \otimes \sigma_k]. \end{aligned} \quad (13)$$

For the case of the generalized four-qubit W-state all vectors $\vec{r}_k$ are collinear, all matrices $g^{(k)}$ are diagonal and the vectors $\vec{r}_k$ are eigenvectors of the matrices $g^{(k)}$ as following:

$$\vec{r}_k = (0, 0, r_k), \quad g_{ij}^{(k)} = \begin{pmatrix} \omega_k & 0 & 0 \\ 0 & \omega_k & 0 \\ 0 & 0 & -\tilde{r}_k \end{pmatrix}, \quad k = 1, 2, 3. \quad (14)$$

In Eq.(14) we defined various quantities as following:

$$\begin{aligned} r_k &= a_1^2 + a_2^2 + a_3^2 + a_4^2 - 2a_k^2, \quad \omega_1 = 2a_2a_3, \quad \omega_2 = 2a_1a_3, \quad \omega_3 = 2a_1a_2. \\ \tilde{r}_1 &= a_2^2 + a_3^2 - a_1^2 - a_4^2, \quad \tilde{r}_2 = a_1^2 + a_3^2 - a_2^2 - a_4^2, \quad \tilde{r}_3 = a_1^2 + a_2^2 - a_3^2 - a_4^2. \end{aligned} \quad (15)$$

In addition, the non-vanishing components of h_{ijk} are

$$h_{113} = h_{223} = \omega_3 \quad h_{131} = h_{232} = \omega_2 \quad h_{311} = h_{322} = \omega_1 \quad h_{333} = -r_4. \quad (16)$$

Due to the maximization in Eq.(12) the Bloch vectors should satisfy the following Lagrange multiplier equations:

$$\begin{aligned} r_{1i} + g_{ij}^{(3)} s_{2j} + g_{ij}^{(2)} s_{3j} + h_{ijk} s_{2j} s_{3k} &= \Lambda_1 s_{1i} \\ r_{2i} + g_{ji}^{(3)} s_{1j} + g_{ij}^{(1)} s_{3j} + h_{kij} s_{1k} s_{3j} &= \Lambda_2 s_{2i} \\ r_{3i} + g_{ji}^{(2)} s_{1j} + g_{ji}^{(1)} s_{2j} + h_{jki} s_{1j} s_{2k} &= \Lambda_3 s_{3i}. \end{aligned} \quad (17)$$

Now we put $s_{1y} = s_{2y} = s_{3y} = 0$ as before. After removing the Lagrange multiplier constants Λ_1 , Λ_2 and Λ_3 , one can show that Eq.(17) reduce to the following three equations:

$$\begin{aligned} s_{1x} [r_1 - \tilde{r}_3 s_{2z} - \tilde{r}_2 s_{3z} + \omega_1 s_{2x} s_{3x} - r_4 s_{2z} s_{3z}] &= s_{1z} [\omega_2 s_{3x} (1 + s_{2z}) + \omega_3 s_{2x} (1 + s_{3z})] \\ s_{2x} [r_2 - \tilde{r}_3 s_{1z} - \tilde{r}_1 s_{3z} + \omega_2 s_{1x} s_{3x} - r_4 s_{1z} s_{3z}] &= s_{2z} [\omega_1 s_{3x} (1 + s_{1z}) + \omega_3 s_{1x} (1 + s_{3z})] \\ s_{3x} [r_3 - \tilde{r}_1 s_{2z} - \tilde{r}_2 s_{1z} + \omega_3 s_{1x} s_{2x} - r_4 s_{1z} s_{2z}] &= s_{3z} [\omega_2 s_{1x} (1 + s_{2z}) + \omega_1 s_{2x} (1 + s_{1z})]. \end{aligned} \quad (18)$$

Eq.(18) implies that the Bloch vectors have the following symmetries:

$$\begin{aligned} \vec{s}_1(a_1, a_2, a_3, a_4) &= \vec{s}_2(a_2, a_1, a_3, a_4) = \vec{s}_3(a_3, a_2, a_1, a_4) \\ \vec{s}_1(a_1, a_2, a_3, a_4) &= \vec{s}_1(a_1, a_3, a_2, a_4) \\ \vec{s}_2(a_1, a_2, a_3, a_4) &= \vec{s}_2(a_3, a_2, a_1, a_4) \\ \vec{s}_3(a_1, a_2, a_3, a_4) &= \vec{s}_3(a_2, a_1, a_3, a_4). \end{aligned} \quad (19)$$

Therefore, one can compute all Bloch vectors if one of them is known. Using the symmetries (19), we can make single equation from Eq.(18) which is expressed in terms of s_{1z} only in a form

$$\frac{s_{1x}(a_1, a_2, a_3, a_4)}{s_{1z}(a_1, a_2, a_3, a_4)} = \frac{P(a_1, a_2, a_3, a_4)}{Q(a_1, a_2, a_3, a_4)} \quad (20)$$

where

$$\begin{aligned}
P(a_1, a_2, a_3, a_4) &= \omega_2 \sqrt{1 - s_{1z}^2(a_3, a_2, a_1, a_4)} [1 + s_{1z}(a_2, a_1, a_3, a_4)] \\
&\quad + \omega_3 \sqrt{1 - s_{1z}^2(a_2, a_1, a_3, a_4)} [1 + s_{1z}(a_3, a_2, a_1, a_4)] \\
Q(a_1, a_2, a_3, a_4) &= r_1 - \tilde{r}_3 s_{1z}(a_2, a_1, a_3, a_4) - \tilde{r}_2 s_{1z}(a_3, a_2, a_1, a_4) \\
&\quad + \omega_1 \sqrt{1 - s_{1z}^2(a_2, a_1, a_3, a_4)} \sqrt{1 - s_{1z}^2(a_3, a_2, a_1, a_4)} \\
&\quad - r_4 s_{1z}(a_2, a_1, a_3, a_4) s_{1z}(a_3, a_2, a_1, a_4).
\end{aligned}$$

Defining $a_1 = a_2 = a_3 \equiv a$ and $a_4 \equiv q$, one can solve Eq.(20) easily. The final expressions of solutions are

$$\begin{aligned}
s_{1z} &= s_{2z} = s_{3z} = \frac{1}{9a^2 - q^2} \\
s_{1x} &= s_{2x} = s_{3x} = \sqrt{1 - s_{1z}^2} = \frac{2\sqrt{6}a}{9a^2 - q^2} \sqrt{3a^2 - q^2}.
\end{aligned} \tag{21}$$

Inserting Eq.(21) into Eq.(12), one can compute P_{max} for $|W_4\rangle$ with $a_1 = a_2 = a_3 \equiv a$ and $a_4 \equiv q$ whose final expression is

$$P_{max} = \frac{2^2(1 - q^2)^3}{(3 - 4q^2)^2}. \tag{22}$$

Eq.(21) implies that P_{max} in Eq.(22) is valid when $q^2 \leq 3a^2$. When $q = 0$, P_{max} becomes $4/9$ as expected. When $q = 1/2$, P_{max} becomes $27/64$, which is in agreement with the result of Ref.[14].

One can repeat the calculation for $|W_5\rangle$ with $a_1 = a_2 = a_3 = a_4 \equiv a$ and $a_5 = q$. Then the final expression of P_{max} becomes

$$P_{max} = \frac{3^3(1 - q^2)^4}{(4 - 5q^2)^3}. \tag{23}$$

When $q = 0$, P_{max} reduces to $27/64$ as expected. When $q = 1/\sqrt{5}$, P_{max} reduces to $(4/5)^4$. By the same way P_{max} for $|W_6\rangle$ can be written as

$$P_{max} = \frac{4^4(1 - q^2)^5}{(5 - 6q^2)^4}. \tag{24}$$

Fig. 1 is a plot of q -dependence of P_{max} for $|W_4\rangle$, $|W_5\rangle$ and $|W_6\rangle$. The black dots are numerical data computed by the numerical technique exploited in Ref.[14]. The red solid and red dotted lines are Eq.(22), Eq.(23) and Eq.(24) when $q \leq 1/\sqrt{2}$ and $q \geq 1/\sqrt{2}$ respectively. As expected the numerical data are in perfect agreement with Eq.(22), Eq.(23) and Eq.(24) in the applicable domain, *i.e.* $q^2 \leq (n-1)a^2$ for $|W_n\rangle$. Outside the applicable domain ($q^2 \geq 1/\sqrt{2}$) the numerical data are in disagreement with these equations.




Fig. 1(a)




Fig. 1(b)




Fig. 1(c)

FIG. 1: Plot of q -dependence of P_{max} for 4-qubit (Fig. 1(a)), 5-qubit (Fig. 1(b)), and 6-qubit (Fig. 1(c)). The black dots are numerical data of P_{max} . The red solid lines are result of Eq.(25) in the applicable domain, $0 \leq q \leq 1/\sqrt{2}$. The red dotted lines are result of Eq.(25) outside the applicable domain. The blue solid lines are plot of $\max(a^2, q^2) = q^2$ outside the applicable domain. This figures strongly suggest that P_{max} for $|W_n\rangle$ is Eq.(25) when $q \leq 1/\sqrt{2}$ and $\max(a^2, q^2) = q^2$ when $q \geq 1/\sqrt{2}$.

V. GENERAL MULTI-QUBIT W-TYPE STATES: ONE-PARAMETRIC CASES

From Eq.(11), (22), (23) and (24) one can guess that P_{max} for W_n is ($a_1 = \dots = a_{n-1} \equiv a$, $a_n \equiv q$)

$$P_{max}(n, q) = (1 - q^2)^{n-1} \left(\frac{n-2}{(n-1) - nq^2} \right)^{n-2}. \quad (25)$$

Using this result, one can straightforwardly construct the nearest product state to $|W_n\rangle$. After some algebra, when $q^2 \leq (n-1)a^2$, one can show that the analytic expression of the

nearest product state is $|q_1\rangle \otimes |q_2\rangle \otimes \cdots \otimes |q_n\rangle$, where

$$\begin{aligned} |q_1\rangle &= \cdots = |q_{n-1}\rangle = \\ &= \frac{1}{\sqrt{(n-1)^2 a^2 - q^2}} \left[\sqrt{(n-1)(n-2)a} |0\rangle + \sqrt{(n-1)a^2 - q^2} e^{i\varphi} |1\rangle \right] \\ |q_n\rangle &= \frac{1}{\sqrt{(n-1)^2 a^2 - q^2}} \left[\sqrt{(n-1)^2 a^2 - (n-1)q^2} |0\rangle + \sqrt{n-2} q e^{i\varphi} |1\rangle \right] \end{aligned} \quad (26)$$

and φ is an arbitrary phase factor. When $q^2 \geq (n-1)a^2$, the nearest product state, of course, becomes $|0 \cdots 01\rangle$.

Now, we present a simple proof for both equations (25) and (26). It is easy to check

$$\langle q_1 q_2 \cdots q_{n-1} | W_n \rangle = e^{-i\varphi} \sqrt{P_{max}} |q_n\rangle, \quad \langle q_2 q_3 \cdots q_{n-1} q_n | W_n \rangle = e^{-i\varphi} \sqrt{P_{max}} |q_1\rangle. \quad (27)$$

The second equation in (27) is invariant under the permutations ($q_1 \leftrightarrow q_j, j = 2, 3, \cdots, n-1$). Thus, the product state satisfies the stationarity equations of Ref.[10] and consequently, is the nearest separable state. Accordingly, $\sqrt{P_{max}}$ is the injective tensor norm of $|W_n\rangle$.

When $q = 0$ and $q = 1/\sqrt{n}$, P_{max} reduces to $(1 - 1/(n-1))^{n-2}$ and $(1 - 1/n)^{n-1}$ respectively. Thus, Eq.(25) is perfectly in agreement with the result of Ref.[14]. Another interesting point in Eq.(25) is that P_{max} becomes $1/2$ regardless of n when $q = 1/\sqrt{2}$, the boundary of the applicable domain. This makes us conjecture that outside the applicable domain P_{max} becomes $\max(a^2, q^2) = q^2$ like 3-qubit case. The blue solid lines in Fig. 1 are plot of q^2 at the domain $q \geq 1/\sqrt{2}$. As we conjecture, the blue lines are perfectly in agreement of numerical data.

Another consequence of Eq.(25) is the entanglement witness $\hat{W}_n$ for an one-parametric W-type state. Its construction is straightforward as following form:

$$\hat{W}_n = P_{max}(n, q) \mathbb{1} - |W_n(q)\rangle \langle W_n(q)|, \quad (28)$$

where $\mathbb{1}$ is a unit matrix. Obviously one can show

$$Tr \left(\hat{W}_n |W_n(q)\rangle \langle W_n(q)| \right) < 0, \quad Tr \left(\hat{W}_n \rho_0 \right) \geq 0, \quad (29)$$

where ρ_0 is any separable state. Thus, $\hat{W}_n$ is an entanglement witness and allows an experimental detection of the multipartite entanglement.

VI. FOUR-QUBIT W STATE: TWO-PARAMETRIC CASES

In this section we will compute P_{max} for the two-parametric $|W_4\rangle$ given by

$$|W_4\rangle = a|1000\rangle + b|0100\rangle + q|0010\rangle + q|0001\rangle. \quad (30)$$

It seems to be difficult to apply the Lagrange multiplier method directly due to their non-trivial nonlinearity. Thus, we will adopt the usual maximization method.

The maximum overlap probability P_{max} is

$$P_{max} = \max_{|q_1\rangle|q_2\rangle|q\rangle} |\langle q_1|\langle q_2|\langle q|W_4\rangle|^2. \quad (31)$$

Now we define the 1-qubit states as $|q_1\rangle = \alpha_0|0\rangle + \alpha_1|1\rangle$, $|q_2\rangle = \beta_0|0\rangle + \beta_1|1\rangle$ and $|q\rangle = \gamma_0|0\rangle + \gamma_1|1\rangle$. For simplicity, we are assuming that all coefficients are real and positive. Then, P_{max} becomes

$$P_{max} = \max_{\alpha_0, \beta_0, \gamma_0} \gamma_0^2 \left(a\beta_0\gamma_0\sqrt{1-\alpha_0^2} + b\alpha_0\gamma_0\sqrt{1-\beta_0^2} + 2q\alpha_0\beta_0\sqrt{1-\gamma_0^2} \right)^2. \quad (32)$$

Since the maximum value is determined at extremum point, it is useful if the extremum conditions are derived. This is achieved by differentiating Eq.(32), which leads to

$$\begin{aligned} b\gamma_0\sqrt{1-\beta_0^2} + 2q\beta_0\sqrt{1-\gamma_0^2} &= a\beta_0\gamma_0\frac{\alpha_0}{\sqrt{1-\alpha_0^2}} \\ a\gamma_0\sqrt{1-\alpha_0^2} + 2q\alpha_0\sqrt{1-\gamma_0^2} &= b\alpha_0\gamma_0\frac{\beta_0}{\sqrt{1-\beta_0^2}} \\ a\beta_0\gamma_0\sqrt{1-\alpha_0^2} + b\alpha_0\gamma_0^2\sqrt{1-\beta_0^2} + q\alpha_0\beta_0\sqrt{1-\gamma_0^2} &= q\alpha_0\beta_0\frac{\gamma_0^2}{1-\gamma_0^2}. \end{aligned} \quad (33)$$

One can solve the equations by separating α_0 from β_0 , γ_0 , i.e.,

$$\begin{aligned} \frac{\alpha_0}{\sqrt{1-\alpha_0^2}} &= \frac{b}{a}\frac{\sqrt{1-\beta_0^2}}{\beta_0} + \frac{2q}{a}\frac{1-\gamma_0^2}{\gamma_0} \\ \frac{\sqrt{1-\alpha_0^2}}{\alpha_0} &= \frac{b}{a}\frac{\beta_0}{\sqrt{1-\beta_0^2}} - \frac{2q}{a}\frac{\sqrt{1-\gamma_0^2}}{\gamma_0} \\ \frac{\sqrt{1-\alpha_0^2}}{\alpha_0} &= \frac{q}{a}\frac{\gamma_0}{\sqrt{1-\gamma_0^2}} - \frac{q}{a}\frac{\sqrt{1-\gamma_0^2}}{\gamma_0} - \frac{b}{a}\frac{\sqrt{1-\beta_0^2}}{\beta_0} \end{aligned} \quad (34)$$

and one can get the solutions for β_0 and γ_0 as follows:

$$\begin{aligned} \beta_0^2 &= \frac{3}{2} - \frac{4q^2 - a^2 + b^2}{4q^2}\gamma_0^2 \\ \gamma_0^2 &= \frac{4q^2(4q^2 - a^2 - b^2) - 2q^2\sqrt{(4q^2 - a^2 - b^2)^2 + 12a^2b^2}}{(4q^2 + b^2 - a^2)^2 - 16q^2b^2}. \end{aligned} \quad (35)$$

The solution for α_0 is obtained by separating β_0 :

$$\alpha_0^2 = \frac{3}{2} - \frac{4q^2 + a^2 - b^2}{4q^2} \gamma_0^2. \quad (36)$$

Inserting these extremum solution in P_{max} and rationalizing denominator, one gets

$$P_{max} = \frac{2q^4 \left[(4q^2 - a^2 - b^2) \{ (4q^2 - a^2 - b^2)^2 - 36a^2b^2 \} + \{ (4q^2 - a^2 - b^2)^2 + 12a^2b^2 \}^{\frac{3}{2}} \right]}{\{ (4q^2 - a^2 - b^2)^2 - 4a^2b^2 \}^2}. \quad (37)$$

Of course, Eq.(37) is valid when $\alpha^2 \leq \beta^2 + \gamma^2 + \delta^2$, where $\{\alpha, \beta, \gamma, \delta\}$ is $\{a, b, q, q\}$ with decreasing order. When $\alpha^2 \geq \beta^2 + \gamma^2 + \delta^2$, P_{max} will be $\alpha^2 = \max(a^2, b^2)$.

The dependence of the maximal overlap on state parameters is shown in Fig.2. The behavior of P_{max} in different limits is explained in the next section.




FIG. 2: The maximal overlap P_{max} vs. the parameters a and b for the 4-qubit state. The green and blue areas are highly entangled regions and the maximal overlap is given by Eq.(37). The violet(dark orange) area is a slightly entangled region and the maximal overlap is $\max(a^2, b^2)$. It is minimal ($P_{max} = 27/64$) at $a = b = 1/2$ which is the W-state and maximal ($P_{max} = 1$) either at $a = 1, b = 0$ or at $a = 0, b = 1$ which are product states.

VII. SPECIAL FOUR-QUBIT W-TYPE STATES

In this section we consider some special 4-qubit states.

The first one is $a = 0$ limit. Since $|W_4\rangle = |0\rangle \otimes (b|100\rangle + q|010\rangle + q|001\rangle)$ in this limit, one can compute P_{max} using Eq.(5). In this limit Eq.(37) gives

$$P_{max} = \frac{4q^4}{4q^2 - b^2} \quad (b^2 \leq 2q^2). \quad (38)$$

One can show easily that this is perfectly in agreement with Eq.(5).

The second special case is $a = q$ limit. In this limit Eq.(37) gives

$$P_{max} = \frac{4(1 - b^2)^3}{(3 - 4b^2)^2} \quad (b^2 \leq 3q^2) \quad (39)$$

which is also consistent with Eq.(22).

The last special case is $2q = a + b$ limit. Although both denominator and numerator in Eq.(37) vanish, their ratio has a finite limit and P_{max} takes correct values in the applicable domain. The applicable domain is defined by the two restrictions $\alpha^2 \leq \beta^2 + \gamma^2 + \delta^2$ and $2q = a + b$. These restrictions together with the normalization condition impose upper and lower bounds for the parameters a and b

$$\min(a, b) \geq \frac{\sqrt{2}}{6}, \quad \max(a, b) \leq \frac{\sqrt{2}}{2}. \quad (40)$$

The maximum overlap probability P_{max} is

$$P_{max} = \frac{27}{256} \frac{(a + b)^4}{ab}. \quad (41)$$

The limit $a = b = q = 1/2$ again yields $P_{max} = 27/64$. Another interesting limit is the case when $b(a)$ is minimal and $a(b)$ is maximal. This limit is reached at $a = 3b$ ($b = 3a$). Then Eq.(41) yields $P_{max} = 1/2 = \alpha^2$. These states are first type shared states[29] and allow perfect teleportation and superdense coding scenario.

VIII. DISCUSSION

We have calculated the maximal overlap of one- and two-parametric W-type states and found their nearest separable states. However, in some sub-region of the parameter space one can find the nearest states and corresponding maximal overlaps for generic W-type states. In fact, the square of any coefficient in Eq.(2) is a maximal overlap in some region of state parameters. It is easy to check that the product state $|0_1 \dots 0_{k-1} 1_k 0_{k+1} \dots 0_n\rangle$ is a solution of

stationarity equation with entanglement eigenvalue $\sqrt{P_{max}} = a_k$. From previous results one can guess that this solution gives a true maximum of the overlap if

$$a_k^2 \geq a_1^2 + a_2^2 + \dots + a_{k-1}^2 + a_{k+1}^2 + \dots + a_n^2 = 1 - a_k^2. \quad (42)$$

Then the maximal overlap in the slightly entangled region can be written readily in the form

$$P_{max} = \max(a_1^2, a_2^2, \dots, a_n^2) \quad \text{if} \quad \max(a_1^2, a_2^2, \dots, a_n^2) \geq \frac{1}{2}. \quad (43)$$

This formula has the following simple interpretation. Equation (42) means that the state is already written in the Schmidt normal form and the maximal overlap takes the value of the largest coefficient [20].

Now the question at issue is what is happening if $a_k^2 < 1/2$, $k = 1, 2, \dots, n$. From these inequalities it follows that

$$\frac{1}{2}(a_1 + a_2 + \dots + a_n) > \max(a_1, a_2, \dots, a_n). \quad (44)$$

From any set of such coefficients one can form polygons(polyhedrons). This fact is an indirect evidence that P_{max} has a geometrical meaning. Unfortunately, there is an obstacle to the goal achievement. The problem is that we have not the answer for generic states. For example, it is difficult to conclude from Eq.(11) that the expression is the circumradius of a triangle in a particular limit. In general, one can form many polygons, either convex or crossed, from the set $a_1, a_2, \dots, a_n$. Each of them generates its own geometric quantities that can be treated as the maximal overlap. This happens because stationarity equations have many solutions in highly entangled region. And all of these solutions yield the same expression in particular cases. For example, in Ref.[29] it was shown that all convex and crossed quadrangles are contracted to the same triangle in particular limits. In conclusion, in order to find a true geometric interpretation one has to derive P_{max} for generic states.

Another(and probably promising) way to get the desired interpretation is the following. Since the surface $(a_1^2 - 1/2)(a_2^2 - 1/2) \dots (a_n^2 - 1/2) = 0$ separates highly and slightly entangled regions, one may ask what is happening on this surface. That is, we are considering polygons whose sides satisfy the equality $a_k^2 = a_1^2 + a_2^2 + \dots + a_{k-1}^2 + a_{k+1}^2 + \dots + a_n^2$ for any k . For $n = 3$ we perfectly know that corresponding polygons are right triangles and the center of a circumcircle lies on the largest side of a right triangle. Then, we can conclude that if the center of the circumcircle is inside the triangle, then the maximal overlap is the circumradius

and otherwise is the largest coefficient. However, for $n \geq 4$ we do not know what are the polygons for which the square of the largest side is the sum of squares of the remaining coefficients. If one understands the geometric meaning of this relation, then one finds a clue. And this clue may enable us to find P_{max} for generic W-type states. These type of analytic expressions can have practical application in QICC and may shed new light on multipartite entanglement.

All above-mentioned problems owe their origin to the fact that the injective tensor norm is related to the Cayley's Hyperdeterminant [23]. It is well-known that this hyperdeterminant has a geometrical interpretation for $n = 3$ and no such interpretation is known for $n \geq 4$ so far. We hope to keep on studying this issue in the future.

Acknowledgments

This work was supported by the Kyungnam University Foundation, 2008.

-
- [1] L. Amico, R. Fazio, A. Osterloh and V. Vedral, *Entanglement in many-body systems*, Rev. Mod. Phys. **80** (2008) 517 [quant-ph/0703044] and references therein.
 - [2] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters, *Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels*, Phys. Rev. Lett. **70**, (1993) 1895.
 - [3] E. Jung, M. R. Hwang, D. K. Park, J. W. Son and S. Tamaryan, *Mixed-state entanglement and quantum teleportation through noisy channels*, J. Phys. A: Math. Theor. **41** (2008) 385302 [arXiv:0804.4595 (quant-ph)].
 - [4] G. Vidal, *Efficient classical simulation of slightly entangled quantum computations*, Phys. Rev. Lett. **91** (2003) 147902 [quant-ph/0301063].
 - [5] R. Jozsa and N. Linden, *On the role of entanglement in quantum computational speed-up*, Proc. R. Soc. Lond. A **459**, (2003) 2011 [quant-ph/0201143].
 - [6] A. K. Ekert, *Quantum cryptography based on Bells theorem*, Phys. Rev. Lett **67**, (1991) 661.

- [7] C. A. Fuchs, N. Gisin, R. B. Griffiths, C. S. Niu and A. Peres, *Optimal eavesdropping in quantum cryptography. I. Information bound and optimal strategy*, Phys. Rev. **A 56**, (1997) 1163 [quant-ph/9701039].
- [8] A. Shimony, *Degree of entanglement*, in D. M. Greenberg and A. Zeilinger (eds.), Fundamental problems in quantum theory: A conference held in honor of J. A. Wheeler, Ann. N. Y. Acad. Sci. **755** (1995) 675.
- [9] H. Barnum and N. Linden, *Monotones and Invariants for Multi-particle Quantum States*, J. Phys. A: Math. Gen. **34**, (2001) 6787 [quant-ph/0103155].
- [10] T.-C. Wei and P. M. Goldbart, *Geometric measure of entanglement and application to bipartite and multipartite quantum states*, Phys. Rev. **A 68** (2003) 042307 [quant-ph/0307219].
- [11] R. Werner and A. Holevo, *Counterexample to an additivity conjecture for output purity of quantum channels*, J. Math. Phys. **43**, 4353(2002).
- [12] L. K. Grover, *Quantum Mechanics helps in searching for a needle in a haystack*, Phys. Rev. Lett. **79** (1997) 325 [quant-ph/9706033].
- [13] O. Biham, M. A. Nielsen and T. J. Osborne, *Entanglement monotone derived from Grover's algorithm*, Phys. Rev. **A 65** (2002) 062312 [quant-ph/0112907].
- [14] Y. Shimoni, D. Shapira and O. Biham, *Characterization of pure quantum states of multiple qubits using the Groverian entanglement measure*, Phys. Rev. **A 69** (2004) 062303 [quant-ph/0309062].
- [15] T.-C. Wei, D. Das, S. Mukhopadhyay, S. Vishveshwara, and P. M. Goldbart, *Global entanglement and quantum criticality in spin chains*, Phys. Rev. **A 71**, (2005) 060305(R) [arXiv:quant-ph/0405162].
- [16] R. Orus, S. Dusuel, and J. Vidal, *Equivalence of critical scaling laws for many-body entanglement in the Lipkin-Meshkov-Glick model*, Phys. Rev. Lett. **101**, (2008) 025701 [arXiv:0803.3151].
- [17] M. Hayashi, D. Markham, M. Mura, M. Owari, and S. Virmani, *Bounds on Multipartite Entangled Orthogonal State Discrimination Using Local Operations and Classical Communication*, Phys. Rev. Lett. **96**, (2006) 040501.
- [18] O. Gühne, M. Reimpell and R.F. Werner, *Estimating entanglement measures in experiments*, Phys. Rev. Lett. **98**, (2007) 110502 [arXiv:quant-ph/0607163].
- [19] H. A. Carteret, A. Higuchi, and A. Sudbery, *Multipartite generalisation of the Schmidt de-*

- composition*, J. Math. Phys. **41**, 7932(2000).
- [20] L. Tamaryan, D. K. Park and S. Tamaryan, *Generalized Schmidt Decomposition based on Injective Tensor Norm* [quant-ph/0809.1290].
 - [21] E. Jung, M. R. Hwang, H. Kim, M. S. Kim, D. K. Park, J. W. Son and S. Tamaryan, *Reduced state uniquely defines Groverian measure of original pure state*, Phys. Rev. **A 77**, (2008) 062317 [quant-ph/0709.4292].
 - [22] T.-C. Wei and S. Severini, *Matrix permanent and quantum entanglement of permutation invariant states*, arXiv:0905.0012v1 [quant-ph].
 - [23] J. J. Hilling and A. Sudbery, *The geometric measure of multipartite entanglement and the singular values of a hypermatrix*, arXiv:0905.2094v2 [quant-ph].
 - [24] S. Tamaryan, T.-C. Wei and D.K. Park, *Maximally entangled three-qubit states via geometric measure of entanglement*, arXiv:0905.3791v1 [quant-ph].
 - [25] R. Hübener, M. Kleinmann, T.-C. Wei, and O. Gühne, *The geometric measure of entanglement for symmetric states*, arXiv:0905.4822v1 [quant-ph].
 - [26] W. Dür, G. Vidal and J. I. Cirac, *Three qubits can be entangled in two inequivalent ways*, Phys.Rev. A **62**, 062314(2000).
 - [27] P. Parashar and S. Rana, *N-Qubit W States are Determined by their Marginals*, arXiv: 0809.4394 [quant-ph].
 - [28] L. Tamaryan, D.K. Park and S. Tamaryan, *Analytic Expressions for Geometric Measure of Three Qubit States*, Phys. Rev. **A 77** (2008) 022325 [quant-ph/0710.0571].
 - [29] L. Tamaryan, D.K. Park, J.-W. Son, S. Tamaryan, *Geometric Measure of Entanglement and Shared Quantum States*, Phys. Rev. **A 78** 032304 [quant-ph/0803.1040].
 - [30] A. Acín, A. Andrianov, L. Costa, E. Jané, J. I. Latorre and R. Tarrach, *Generalized Schmidt Decomposition and Classification of Three-Quantum-Bit States*, Phys. Rev. Lett. **85** (2000) 1560 [quant-ph/0003050].
 - [31] E. Jung, M. R. Hwang, D. K. Park, L. Tamaryan and S. Tamaryan, *Three-Qubit Groverian Measure*, Quant. Inf. Comp. **8** (No. 10) (2008) 0925 [quant-ph/0803.3311].